

Instructions for Use
GONITRO™ (GO-NYE-troh)
(nitroglycerin) sublingual powder

Read these Instructions for Use before you start taking GONITRO and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or your treatment.

Important information:

- **GONITRO is for use under the tongue.**
- A dose of GONITRO may be either 1 or 2 packets. Follow your healthcare provider's instructions about how many packets you should take for each dose. After you take your first dose of GONITRO, you may take 1 additional packet of GONITRO every 5 minutes, if needed.
- **You should not take more than 3 packets of GONITRO within 15 minutes.**
- **Get emergency medical help right away if you still have chest pain after taking a total of 3 packets of GONITRO.**
- It is best to take GONITRO while you are resting and in a sitting position.

Taking a dose of GONITRO

Step 1.

Hold the packet of GONITRO upright with the notch and red arrow line at the top of the packet.

Step 2.

Tap the bottom of the GONITRO packet so that the powder settles at the bottom of the packet.

Step 3.

Hold the GONITRO packet at the notch. Hold the packet as close to your mouth as possible and tear along the red arrow line.

Step 4.

Lift up your tongue.

Step 5.

Pour all of the powder in the GONITRO packet under your tongue.

Step 6.

Close your mouth right away and breathe normally through your nose. Allow all of the powder to dissolve before you swallow. Do not rinse your mouth or spit for 5 minutes after taking GONITRO.

Step 7.

If another dose of GONITRO is needed, repeat Steps 1 through 6.

Regularly check the number of packets remaining in your GONITRO box and refill your prescription when there are three packets remaining.

How should I store GONITRO?

Store GONITRO at room temperature between 68°F to 77°F (20°C to 25°C).

Keep GONITRO and all medicines out of the reach of children.

These Instructions for Use have been approved by the U.S. Food and Drug Administration.

Manufactured for
Espero Pharmaceuticals, Inc.
Jacksonville, FL 32250 (800) 235-6520


PHARMACEUTICALS

by G. Pohl-Boskamp GmbH & Co. KG 25551 Hohenlockstedt, Germany.

